



UNIVERSITAS WIRARAJA

LEMBAGA PENELITIAN DAN PENGABDIAN KEPADA MASYARAKAT

Kampus : Jl. Raya Sumenep Pamekasan KM. 5 Patean, Sumenep, Madura 69451 Telp : (0328) 664272/673088
e-mail : lppm@wiraraja.ac.id Website : lppm.wiraraja.ac.id

SURAT PERNYATAAN

Nomor : 200/SP.HCP/LPPM/UNIJA/XII/2020

Yang bertanda tangan di bawah ini :

Nama : Anik Anekawati, M.Si
Jabatan : Kepala LPPM
Instansi : Universitas Wiraraja

Menyatakan bahwa :

Nama : Syaifurrahman Hidayat, S.Kep., Ns., M.Kep.
Jabatan : Staf Pengajar Fakultas Ilmu Kesehatan

Telah melakukan cek plagiarisme ke LPPM menggunakan *software turnitin.com* untuk artikel dengan judul "*Risk Factors for Low Birth Weight In Puskesmas Ganding*" dan mendapatkan hasil similarity sebesar 13%

Demikian surat pernyataan ini dibuat untuk digunakan dengan sebaik-baiknya.

Sumenep, 08 Desember 2020

Kepala LPPM
Universitas Wiraraja,

Anik Anekawati, M.Si
NIDN. 0714077402

RISK FACTORS FOR LOW BIRTH WEIGHT IN PUSKESMAS GANDING

by Syaifurrahman Hidayat

Submission date: 03-Dec-2020 03:26PM (UTC+0700)

Submission ID: 1463372901

File name: EJMCM_Volume_7_Issue_7_Pages_21-27.pdf (245.21K)

Word count: 3355

Character count: 17774

RISK FACTORS FOR LOW BIRTH WEIGHT IN PUSKESMAS GANDING

Ratna Indriyani¹, Ahmaniyah², Mujib Hannan³, Syaifurrahman Hidayat⁴

e-mail:ratnaindriyani@wiraraja.ac.id,²ahmaniyah.fik@wiraraja.ac.id
³mujib@wiraraja.ac.id⁴dayat.fik@wiraraja.ac.id

^{1,2}Lecturer of Midwifery Study Program of Wiraraja University

^{3,4}Lecturer of Nursing Study Program of Wiraraja University

ABSTRACT: Background: The high incidence of low birth weights (LBW) is due to interrelated factors from both the mother and the fetus. Maternal health status affects birth weight directly. Periodic health check for pregnant women (antenatal care visits) are one of the main activities in maintaining the health of the mother and the fetus.

Objective: To examine the risk factors for LBW.

Method: This was an observational analytical study with a case-control approach. The sample was 48 respondents, with 24 mothers experiencing LBW as the case respondents and 24 mothers whose babies were in the normal weight range as the control respondents. The size of the sample was determined using a 1:1 comparison with a purposive sampling method.

Findings: The average height of the respondents was 151.23 cm, weight gain was 9.00 kg, systolic blood pressure was 113.75, mid-upper arm circumference (MUAC) was 23.58 cm, gestational age was 37.42 weeks, and infant birth weight was 2604.17 g. On average, the examination of multiparous status, fetal position, head presentation, counseling, and laboratory tests were also determined. A chi-squared test showed no statistically significant relationships with LBW for the height of the pregnant woman, the blood pressure, the fetal position, the status of TT vaccinations, the administration of iron tablets, counseling, or laboratory examination. In contrast, weight gain, MUAC, fundal height, parity, and maternal age had significant relationships with LBW.

Conclusion: The maternal age, weight gain during pregnancy, MUAC, fundal height, and parity were associated with LBW.

Keywords: LBW, antenatal care

1. BACKGROUND

The World Health Organization (WHO) defines low birth weight (LBW) babies as those born with a bodyweight below 2500 gram. LBW is a major indicator of the health and nutrition of the mother and the fetus. Newborns with LBW have a higher risk of death in the first 28 days of life.^{1,2}

More than 20 million babies were born with LBW or around 1 in 7 births in the world, according to 2015 data. This means in a decrease in LBW in 2000 and 2015 by 1.2%, and WHO has targeted a 30% reduction in the prevalence of LBW between 2012 and 2025. Thus, a 2.7% reduction in LBW incidence rates each year is needed to meet the target.³

The results of the RISKESDAS survey in Indonesia confirmed an increase of LBW by 5.4 to 6.2.4 between 2007 to 2018, and infant and neonatal mortality rates for LBW babies was 47%. The main causes of death were premature birth, asphyxia, infection, and birth defects.⁵ LBW is the result of preterm labor (gestational age < 37 weeks)—mothers who give birth before a gestational age of 37 weeks are ten times more likely to give birth to LBW babies compared with mothers having a full-term birth (≥ 37 weeks). Increased LBW is the main predictor of infant mortality, especially in the first month of life, and those who survive are more likely to suffer growth retardation, and lower IQ⁶ and its effects will continue into adulthood, increasing the risk of chronic conditions such as obesity and diabetes.⁷

Many interrelated factors (both from the mother and the fetus) increase the risk of LBW. Maternal health status affects birth weight directly. Periodic health check-ups on pregnant women (antenatal care visits) are one of the main activities for maintaining the health of the mother and the fetus. The importance of antenatal care for pregnant women is related to the promotion of health, screening and diagnosis, and the prevention of disease. WHO recommends that antenatal care should consist of at least four visits for a woman with a normal pregnancy.⁸ Regular antenatal care is the first step in preventing LBW.⁹ Consequently, the quality and quantity of antenatal care must be maintained. Antenatal care in Indonesia is carried out in an integrated manner comprising ten examination measures: measurement of height and weight, measurement of blood pressure, measurement of fundal height, measurement of mid-upper arm circumference (MUAC), examination of the fetal position and fetal heart rate, a check on the status of the tetanus toxoid (TT) vaccination, administration of blood-booster tablets, laboratory tests, pregnancy counseling, and counseling related to health problems during pregnancy. These services must be available for pregnant women since these factors may contribute to the risk of LBW. After consideration of these measures, we decided to study the risk factors of LBW in *Puskesmas Ganding* to determine which risk factors were associated with LBW.

2. METHOD

This was an observational analytical study with a case-control approach conducted at *Puskesmas Ganding* of Sumenep Regency from January to December 2019. Our study involved 48 mothers divided into two groups: 24 mothers experiencing LBW as the case respondents, and 24 mothers with normal weight babies as the control respondents. The size of the sample was determined using a 1:1 comparison with a purposive sampling method.

The independent variables were the ten antenatal care measures taken by midwives (measurement of height and weight, measurement of blood pressure, measurement of fundal height, measurement of MUAC, examination of the fetal position and fetal heart rate, a check on the status of the tetanus toxoid vaccination, administration of blood-booster tablets, laboratory tests, pregnancy counseling, and counseling related to health problems during pregnancy) documented in the KIA book. The dependent variable was LBW. We used a bivariate analysis with a chi-squared test to determine which of the independent variables were associated with LBW.

3. FINDINGS AND DISCUSSION

The average height of the respondents was 151.23 cm, the average weight gain was 9.00 kg, systolic blood pressure was 113.75, MUAC was 23.58 cm, gestational age was 37.42 weeks, and infant birth weight was 2604.17 g. The parity of multiparous children, fetal position, head presentation, counseling, and laboratory tests were also conducted.

Those factors are the rights of pregnant women in the integrated ANC examination at *Puskesmas Ganding*. A bivariate analysis was done after the examination to reveal the factors affecting LBW at *Puskesmas Ganding*, see Table1.

Table 1. LBW Risk Factors

Variable	Case Group	Control Group	P-Value
	n %	n %	
Maternal age			
< 20 years old	3 (12.5%)	5 (20.83%)	0.05
20 – 35 years old	8(33.33%)	13 (54.16%)	
>35 years old	13 (54.16%)	6 (25%)	
Height			
< 145 cm	2 (8.33%)	1 (4.16%)	0.80
>145 cm	22(91.66%)	23 (95.83%)	
Weight			
< 9 kg	22 (91.22%)	0	0.00
> 9 kg	2 (8.33%)	24 (100%)	
Blood pressure (systolic)			
< 120	10 (41.66%)	12 (50%)	0.24
≥ 120	14 (58.33%)	12 (50%)	
MUAC			
< 23.5 cm	16 (66.66%)	1(4.16%)	0.00
≥ 23.5cm	8 (33.33%)	23 (95.83%)	
Fundal Height			
Suitable	6 (25%)	24 (100%)	0.00
Not suitable	18 (75%)	0	
Fetal Position			
Vertex presentation	23 (95.83%)	22 (91.66%)	0.20
Breech presentation	1 (4.16)	2 (8.33%)	
TT Vaccination			
Complete (5TT)	24 (100%)	24 (100%)	0.23
Not complete	0	0	
Administration of Fe tablet			
Yes	24 (100%)	24 (100%)	0.23
No	0	0	
Pregnancy Counseling			
Yes	24 (100%)	24 (100%)	0.08
No	0	0	
Parity			
Primiparous	3 (12.5%)	12 (50%)	0.00
Multiparous	7 (29.16%)	9 (37.5%)	
Grand multiparous	14 (58.33%)	3 (12.5%)	
Laboratory checks			
Yes	24 (100%)	24 (100%)	0.33
No	0	0	

The chi-square test showed no significant relationship with LBW for the height of the pregnant woman ($p = 0.080$), the blood pressure ($p = 0.24$), the fetal position ($p = 0.20$), the

status of the TT vaccination ($p = 0.20$), the administration of iron tablets ($p = 0.21$), counseling ($p = 0.08$), or laboratory checks ($p = 0.33$) because the value of $p > 0.05$. In contrast, weight gain ($p = 0.00$), MUAC ($p = 0.00$), fundal height ($p = 0.00$), parity (0.00), and maternal age ($p = 0.05$) showed significant relationships with LBW because the value of $p \leq 0.05$.

The relationship between height and LBW was not significant because most mothers in the case and control groups had an average height of more than 145 cm—a study by Ismi confirmed that heights > 145 cm were not a risk factor for LBW.¹⁰ In contrast, mothers' height also has a relationship with stunting. Short mothers with a height of < 145 cm have a 2.14-times higher risk of having children experiencing stunting than mothers of a normal height.¹¹

2 There was no significant relationship between blood pressure and LBW because mothers in both groups had normal blood pressure (below 130 mmHg), which means that they did not suffer from hypertension. Hypertension is a triad of preeclampsia that causes LBW. Thus, for blood pressure, the risk of LBW can be predicted with a sensitivity and specificity of 70%.¹²

The fetal position had no significant relationship with LBW because most of the fetuses had a normal or vertex presentation. Normal presentation and head presentation all determine the fetal position, which affects the delivery process (normal or abnormal). Our study confirmed that the delivery process had no significant relationship with LBW¹³, and theoretically, there has been no pathophysiology as well.

2 There was no significant relationship between TT vaccination and LBW, as most of the respondents had been vaccinated. TT is a protein-based subunit vaccine that elicits IgG1 immune responses, with antibodies actively transported across the placenta with efficiencies $> 100\%$. WHO recommends giving two doses of TT to be given twice, one in the first pregnancy and one for every subsequent pregnancy for a maximum of five doses.¹⁴ Pregnant women who have not had a TT vaccination have 9.50 times more risk of experiencing LBW than those who have had the vaccination.¹⁵

Iron administration had no significant relationship with LBW because all the respondents had taken iron tablets. However, there were LBW cases in the case group, and some respondents were not taking their iron tablets as recommended. Pregnant women who consume fewer than 90 iron tablets have a 2.1 times greater risk of LBW compared with those who consume more than 90 iron tablets.¹⁶ The non-compliance in consuming iron tablets leads to pregnant women suffering from anemia, which has a very significant impact on LBW.^{17, 18} Our study did not measure the number of iron tablets given during pregnancy, so the statistical test showed no significant relationship between the administration of iron tablets and LBW.

Pregnancy counseling and counseling related to health problems during pregnancy also had no significant relationship with LBW. All of the respondents visiting midwives were given the choice of pregnancy counseling from health workers. However, there was a shortage of time for counseling, and mothers could not gain a comprehensive knowledge and understanding of their pregnancy. This might have contributed to the LBW in the case group. Good counseling results in a comfortable and warm relationship between mothers and health workers that can help to create certainty for the mothers.¹⁹

1 There was no significant relationship between LBW and the laboratory tests and examinations. Every pregnant mother visiting *Puskesmas Ganding* had to have a laboratory

examination to assess the health status of the mother and the fetus. Although laboratory tests were conducted, LBW still occurred—thus, further investigations are needed because laboratory tests at *Puskesmas Ganding* only covered hematological examinations (checking hemoglobin and blood type). Blood chemistry and complete urine examinations were only done if mothers mentioned that they were suffering from a complaint. We believe that complete hematological, blood chemistry, and urine examinations could help to predict LBW.

1 There was a significant relationship between the independent variables – weight gain, MUAC, fundal height, parity, and gestational age – and the dependent variable LBW. Weight gain during pregnancy in the case group was mostly below 9 kg as recommended by WHO, who recommended that weight gain in pregnancy should be around 9 kg with a weight gain of 1 kg per month. The baby's birth weight is affected by the baby's weight gain during pregnancy, and maternal weight gain must be in harmony with the fetus's weight gain because a lack of nutrition in mothers indirectly leads to LBW and affects the delivery process.^{20,21} MUAC had a significant relationship with LBW because it represents the nutritional status of the pregnant women. If mothers experience a lack of nutrition during pregnancy, there is a higher possibility of LBW, so mothers must get counseling to understand their nutritional needs during pregnancy.²² This finding was consistent with the results of similar studies.^{23, 24} Fundal height had a significant relationship with LBW since it helps to predict a baby's weight and gestational age with a high degree of accuracy before delivery.²⁵ Measurement of fundal height is a mandatory procedure in antenatal care. Parity was also related to LBW events because the average parity of the respondents was > 3 and represented the nutritional status of mothers. Parities > 3 represent a 1.9 times risk of LBW compared with parities < 3.²⁶ Maternal age also had a significant relationship with LBW because of the decrease in the function reproductive organs when women reach the age of 35 or more. A higher risk for various complications also occurs in pregnant women over the age of 35. Studies of older pregnant women showed that they were more vulnerable to preeclampsia, hypertension related to pregnancy, and cesarean sections.²⁷ Our findings supported other studies which found that maternal age affected infant birth weight or LBW.²⁸⁻³⁰ Weight gain, MUAC, fundal height, parity, and gestational age are related to each other, and all of them can be used to predict LBW if the antenatal examination is performed well. Thus, the quality of antenatal care needs to be improved.

4. CONCLUSIONS

This study investigated ten factors that could influence LBW. We found that LBW was affected by maternal age, weight gain during pregnancy, mid-upper arm circumference, fundal height, and parity. A detailed assessment and examination are needed to improve the quality of antenatal care so that LBW can be minimized.

Cohort studies to determine the factors responsible for LBW are recommended as a follow-up of the present study.

5. REFERENCES

- [1] UNICEF-WHO. UNICEF-WHOLOW BIRTHWEIGHT ESTIMATES Levels and trends 2000–2015 [cited 2020 14 April]. Available from: <https://www.unicef.org/media/53711/file/UNICEF-WHO%20Low%20birthweight%20estimates%202019%20.pdf>.
- [2] WHO. Low-birth-weight newborns (%) 2020 [cited 2020 18 April]. Available from: <https://www.who.int/data/gho/indicator-metadata-registry/imr-details/76>.

- [3] UNICEF-WHO. 1 in 7 babies worldwide born with a low birth weight - The Lancet Global Health, UNICEF, WHO [cited 2020 14 April]. Available from: <https://www.unicef.org/press-releases/1-7-babies-worldwide-born-low-birthweight-lancet-global-health-unicef-who>.
- [4] RISKESDAS. hasil utama Riskesdas 2018 2019 [cited 2020 6 April]. Available from: http://kesmas.kemkes.go.id/assets/upload/dir_519d41d8cd98f00/files/Hasil-riskesdas-2018_1274.pdf.
- [5] Achadi EL. kematian maternal dan neonatal di indonesia 2019. Available from: <https://www.kemkes.go.id/resources/download/info-terkini/rakerkesnas-2019/SESI%20I/Kelompok%20I/1-Kematian-Maternal-dan-Neonatal-di-Indonesia.pdf>.
- [6] Gu H, Wang L, Liu L, Luo X, Wang J, Hou F, et al. A gradient relationship between low birth weight and IQ: A meta-analysis. *Scientific reports*. 2017;7(1):1-13.
- [7] Jornayvaz FR, Vollenweider P, Bochud M, Mooser V, Waeber G, Marques-Vidal P. Low birth weight leads to obesity, diabetes and increased leptin levels in adults: the CoLaus study. *Cardiovascular diabetology*. 2016;15(1):73.
- [8] WHO. WHO recommendations on antenatal care for a positive pregnancy experience: World Health Organization; 2016.
- [9] Ahmed Z, Khoja S, Tirmizi SS. Antenatal care and the occurrence of Low Birth Weight delivery among women in remote mountainous region of Chitral, Pakistan. *Pakistan Journal of Medical Sciences*. 2012;28(5):800.
- [10] Ismi Trihardiani IT. Faktor risiko kejadian berat badan lahir rendah di wilayah kerja Puskesmas Singkawang Timur dan Utara Kota Singkawang: Diponegoro University; 2011.
- [11] Nasution D, Nurdiati DS, Huriyati E. Berat badan lahir rendah (BBLR) dengan kejadian stunting pada anak usia 6-24 bulan. *jurnal gizi klinik Indonesia*. 2014;11(1):31-7.
- [12] Yadav H, Lee N. Maternal factors in predicting low birth weight babies. *The Medical journal of Malaysia*. 2013;68(1):44-7.
- [13] Durie DE, Sciscione AC, Hoffman MK, Mackley AB, Paul DA. Mode of delivery and outcomes in very low-birth-weight infants in the vertex presentation. *American journal of perinatology*. 2011;28(03):195-200.
- [14] Chu HY, Englund JA. Maternal immunization. *Clinical infectious diseases*. 2014;59(4):560-8.
- [15] Aida F. Analisis Faktor Determinan Bayi Berat Badan Lahir Rendah (BBLR) di Kabupaten Sijunjung Tahun 2016: Universitas Andalas; 2017.
- [16] Fatimatasari F, Hadi H, Rahmawati NI. Kepatuhan Mengonsumsi Tablet Fe Selama Hamil Berhubungan dengan Kejadian Bayi Berat Lahir Rendah (BBLR) di Kabupaten Bantul. *Jurnal Ners dan Kebidanan Indonesia*. 2013;1(3):87-9.
- [17] Riu D, Mappaware N, Asmi M, Tansil A. Maternal hemoglobin concentration and birth weight: A report from mother and child tertiary hospital. *Enfermería Clínica*. 2020;30:92-5.
- [18] Iqbal S, Ekmekcioglu C. Maternal and neonatal outcomes related to iron supplementation or iron status: a summary of meta-analyses. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2019;32(9):1528-40.
- [19] Nicoloro-SantaBarbara J, Rosenthal L, Auerbach MV, Kocis C, Busso C, Lobel M. Patient-provider communication, maternal anxiety, and self-care in pregnancy. *Social Science & Medicine*. 2017;190:133-40.
- [20] Rijvi S, Abbasi S, Karmakar A, Siddiqua SF, Dewan F. A Study on Maternal Weight Gain and its Correlation with Birth Weight of Baby at Term. *Anwer Khan Modern Medical College Journal*. 2018;9(1):22-8.

- [21] Trisnawati Y, Utami T. Hubungan Kenaikan Berat Badan Ibu Selama Hamil dengan Kejadian Berat Badan Lahir Rendah (BBLR) di RSUD Kota Tanjungpinang Tahun 2017. *Cakrawala Kesehatan: Kumpulan Jurnal Kesehatan*. 2017;8(1).
- [22] Siyoum M, Melese T. Factors associated with low birth weight among babies born at Hawassa University Comprehensive Specialized Hospital, Hawassa, Ethiopia. *Italian journal of pediatrics*. 2019;45(1):48.
- [23] Nurhayati N, Hamang SH, Thamrin H. Faktor Risiko Umur, Paritas, dan LiLA Ibu Hamil terhadap Kejadian BBLR di RSKDIA Siti Fatimah Makassar. *Window of Midwifery Journal*. 2020:33-41.
- [24] Kusparlina EP. Hubungan antara umur dan status gizi ibu berdasarkan ukuran lingkaran atas dengan jenis BBLR. *Jurnal Penelitian Kesehatan " SUARA FORIKES"(Journal of Health Research" Forikes Voice)*. 2016;7(1).
- [25] Lee AC, Whelan R, Bably NN, Schaeffer LE, Rahman S, Ahmed S, et al. Prediction of gestational age with symphysis-fundal height and estimated uterine volume in a pregnancy cohort in Sylhet, Bangladesh. *BMJ open*. 2020;10(3):e034942.
- [26] Lisdiana AB, Hidayat A. FAKTOR IBU YANG BERHUBUNGAN DENGAN KEJADIAN BERAT BADAN LAHIR RENDAH (BBLR) DI RSUD WONOSARI GUNUNGKIDUL YOGYAKARTA: Universitas' Aisyiyah Yogyakarta; 2017.
- [27] Tseng K-T, Peng C-C, Chang J-H, Hsu C-H, Lin C-Y, Jim W-T, et al. The impact of advanced maternal age on the outcomes of very low birth weight preterm infants. *Medicine*. 2019;98(5).
- [28] Suwarni Y, Noor MS, Rahayu A. Hubungan antara Paritas, LILA, Kadar Hb dan Usia Ibu Hamil dengan Berat Lahir Bayi. *Jurnal Publikasi Kesehatan Masyarakat Indonesia*. 2016;1(1).
- [29] Pamungkas RS. Hubungan Usia Ibu dan Paritas dengan Tingkat Kejadian BBLR di Wilayah Kerja Puskesmas Plered, Kecamatan Plered Kabupaten Purwakarta Tahun 2014. 2015.
- [30] Kim H-E, Song IG, Chung S-H, Choi Y-S, Bae C-W. Trends in Birth Weight and the Incidence of Low Birth Weight and Advanced Maternal Age in Korea between 1993 and 2016. *Journal of Korean medical science*. 2019;34(4).

RISK FACTORS FOR LOW BIRTH WEIGHT IN PUSKESMAS GANDING

ORIGINALITY REPORT

13%

SIMILARITY INDEX

11%

INTERNET SOURCES

7%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	scifes.fkm.ui.ac.id Internet Source	2%
2	repo.unand.ac.id Internet Source	2%
3	"1st Annual Conference of Midwifery", Walter de Gruyter GmbH, 2020 Publication	1%
4	Genevieve K. Dudar, Luca D. D'Andrea, Rossella Di Stasi, Carlo Pedone, John L. Wallace. "A vascular endothelial growth factor mimetic accelerates gastric ulcer healing in an iNOS-dependent manner", American Journal of Physiology-Gastrointestinal and Liver Physiology, 2008 Publication	1%
5	www.unicef.org Internet Source	1%
6	periodicos.unifor.br Internet Source	1%

7	pjms.com.pk Internet Source	1%
8	academic.oup.com Internet Source	1%
9	Shanshan Wang, Liren Yang, Li Shang, Wenfang Yang, Cuifang Qi, Liyan Huang, Guilan Xie, Ruiqi Wang, Mei Chun Chung. "Changing trends of birth weight with maternal age: a cross-sectional study in Xi'an city of Northwestern China", Research Square, 2020 Publication	1%
10	elibrary.almaata.ac.id Internet Source	1%
11	Submitted to University of KwaZulu-Natal Student Paper	1%
12	Chu, H. Y., and J. A. Englund. "Maternal Immunization", Clinical Infectious Diseases, 2014. Publication	1%
13	www.panafrican-med-journal.com Internet Source	1%
14	www.iosrjournals.org Internet Source	<1%
15	imu.edu.my Internet Source	<1%

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On